arXiv:1601.04480v3 [math.GR] 6 Sep 2016

ONE-RELATOR MAXIMAL PRO-p GALOIS GROUPS AND KOSZUL ALGEBRAS

C. QUADRELLI

To John P. Labute, with respect and admiration.

Abstract. Let p be a prime number and let K be a field containing a root of 1 of order p. If the maximal pro-p Galois group $G_K(p)$ is a finitely generated one-relator pro-p group, we show that L. Positselski's and T. Weigel's Koszulity conjectures are true for K. Also, under the above hypothesis we show that the $\mathbb{F}_p$-cohomology algebra of $G_K(p)$ is the quadratic dual of a canonical graded algebra induced by the complete group algebra $\mathbb{F}_p[\![G_K(p)]\!]$, which decomposes as free product of quadratic algebras, in analogy with a result by T. Würfel on one-relator absolute pro-p Galois groups.

1. Introduction

For a prime number p and a field K, let $K(p)$ be the maximal p-extension of K — namely, $K(p)$ is the compositum of all finite Galois p-extensions L/K. The Galois group of $K(p)/K$, called the *maximal pro-p Galois group* of K and denoted by $G_K(p)$, is the maximal pro-p quotient of the absolute Galois group G_K of K (note that the class of maximal pro-p Galois groups includes also absolute Galois groups which are pro-p). Characterizing pro-p groups which are realizable as maximal pro-p Galois groups is one of the major problems in current research in Galois theory.

A pro-p group G is said to be a *one-relator* pro-p group if it has a minimal presentation with only one defining relation. I.e., $G = F/R$, with F a free pro-p group (such that G and F have the same minimal number of generators) and R is generated as closed normal subgroup of F by one element. In [Wür85] T. Würfel proved the following characterization for absolute Galois groups which are one-relator pro-p groups: if such G_K is finitely generated and K contains all the roots of unity of order a p-power, then G_K is a free-by-Demushkin group — i.e., one may find a free pro-p normal subgroup N of G_K such that the quotient G_K/N is a Demushkin group. (Demushkin groups are Poincaré pro-p groups of cohomological dimension two, and they appear as maximal pro-p Galois groups of certain local fields, see Section 2.3).

In order to obtain such result, Würfel used the Merkurjev-Suslin theorem (cf. [MS82]), which is the "degree-2 step" of the well-known Bloch-Kato conjecture, proved by M. Rost and V. Voevodsky (cf. [NSW08, Thm. 6.4.3]). In particular,

Date: September 7, 2016.

2010 *Mathematics Subject Classification.* Primary 12G05; Secondary 17A45, 20F40, 20F14.

Key words and phrases. Galois cohomology, Koszulity conjecture, one-relator pro-p groups, quadratic algebras, Demushkin groups.

The author is supported by the Israel Science Fundation grant no. 152/13.

the Bloch-Kato conjecture implies that if a field K contains a root of unity of order p, then the $\mathbb{F}_p$-cohomology ring $H^\bullet(G_K(p),\mathbb{F}_p) = \bigoplus_{n\geq 0} H^n(G_K(p),\mathbb{F}_p)$, endowed with the (graded-commutative) cup product

$$\cup\colon H^r(G_K(p),\mathbb{F}_p)\times H^s(G_K(p),\mathbb{F}_p)\longrightarrow H^{r+s}(G_K(p),\mathbb{F}_p),$$

is a *quadratic $\mathbb{F}_p$-algebra*, i.e., such algebra is generated by elements of degree one and its defining relations are generated by (homogeneous) relations of degree two (cf. [Qua14, § 2]). Our study of maximal pro-p Galois group which are finitely generated one-relator pro-p groups starts from the following observation.

Observation 1.1. *Let K be a field containing a root of unity of order p (and also $\sqrt{-1}$ if $p=2$), and assume that the maximal pro-p Galois group $G_K(p)$ is a finitely generated one-relator pro-p group. Then one has the decomposition as free product of quadratic $\mathbb{F}_p$-algebras*

$$H^\bullet(G_K(p),\mathbb{F}_p)\simeq \big(\mathbb{F}_p\oplus V_1\oplus H^2(G_K(p),\mathbb{F}_p)\big)\sqcap(\mathbb{F}_p\oplus V_2), \tag{1.1}$$

with $V_1\oplus V_2 = H^1(G_K(p),\mathbb{F}_p)$, where the cup product induces a non-degenerate pairing $V_1\times V_1\to H^2(G_K(p),\mathbb{F}_p)$, and $V_2 = H^1(G_K(p),\mathbb{F}_p)^\perp$.

For a quadratic $\Bbbk$-algebra $A_\bullet = \bigoplus_{n\geq 0} A_n$ one defines the *quadratic dual* $A^!_\bullet$ as the quadratic $\Bbbk$-algebra generated by the dual space A_1^* of A_1, and whose relations are the orthogonal space of the generating relations of $A_\bullet$ (see [PP05, § 1.2] and Definition 2.1). It is natural to ask what is the quadratic dual of the $\mathbb{F}_p$-cohomology ring of a maximal pro-p Galois group: for maximal pro-p Galois groups which are finitely generated one-relator pro-p groups we prove the following.

Theorem 1.2. *Let K be a field containing a root of unity of order p such that $G_K(p)$ is a finitely generated one-relator pro-p group. The quadratic dual of the $\mathbb{F}_p$-cohomology ring of $G_K(p)$ is the graded group algebra*

$$\mathrm{gr}_\bullet(\mathbb{F}_p[\![G_K(p)]\!]) = \bigoplus_{n\geq 0} I^n/I^{n+1}, \qquad \text{with } I^0 = \mathbb{F}_p[\![G_K(p)]\!],$$

where I is the augmentation ideal of the complete group algebra $\mathbb{F}_p[\![G_K(p)]\!]$. Also, such graded algebra splits as free product

$$\mathrm{gr}_\bullet(\mathbb{F}_p[\![G_K(p)]\!])\simeq D_\bullet\sqcup\mathbb{F}_p\langle X\rangle, \tag{1.2}$$

where $D_\bullet$ is a quadratic Demushkin algebra and $\mathbb{F}_p\langle X\rangle$ is the free (graded) associative algebra on a set of indeterminates $X=\{X_1,\dots,X_m\}$.

Demushkin algebras are one-relator (non-commutative) polynomial algebras related to Demushkin groups (to be defined in Section 2.3). In [MPQT] it is conjectured that the graded group algebra $\mathrm{gr}_\bullet(\mathbb{F}_p[\![G_K(p)]\!])$ is the quadratic dual of the $\mathbb{F}_p$-cohomology ring $H^\bullet(G_K(p),\mathbb{F}_p)$ for every field K containing a root of unity of order p such that $G_K(p)$ is finitely generated. Thus, Theorem 1.2 provides a positive answer in the case of one-relator maximal pro-p Galois groups.

Note that by (1.1) and (1.2) the two algebras $H^\bullet(G_K(p),\mathbb{F}_p)$ and $\mathrm{gr}_\bullet(\mathbb{F}_p[\![G_K(p)]\!])$ split in a "Demushkin part" and in a "free part". In fact, we prove a result for maximal pro-p Galois groups which is analogous to Würfel's one (cf. Theorem 5.1).

For a graded $\Bbbk$-algebra $A_\bullet$, one defines the $\Bbbk$-cohomology of $A_\bullet$ as the derived functor of $\mathrm{Hom}_{A_\bullet}(\Bbbk,\Bbbk)$, which is endowed with an additional internal grading —

induced by the grading of $A_\bullet$ —, so that one gets a bigraded algebra. If the non-trivial cohomology groups are concentrated on the diagonal, $A_\bullet$ is called a *Koszul* $\Bbbk$-*algebra* (cf. [PP05, § 1.1, § 2.1]). Koszul algebras are rather mysterious. They were introduced by S. Priddy in [Pri70], and they arise in various fields of mathematics, such as representation theory, noncommutative algebra, noncommutative geometry, algebraic geometry and topology.

Koszul algebras have been studied in the context of Galois cohomology by L. Positselski and A. Vishik (cf. [PV95]). In particular, Positselski conjectured that if a field K contains a root of unity of order p, then the $\mathbb{F}_p$-cohomology algebra of $G_K(p)$ is Koszul, and he showed that such conjecture is a generalization of the Bloch-Kato conjecture (cf. [Pos05, Pos14]). More recently, Koszul algebras have been studied in the context of Lie algebras and pro-p groups by T. Weigel (cf. [Wei15a, Wei15b]). In particular, Weigel conjectured that under the same assumptions on the field K, the graded group algebra $\mathrm{gr}_\bullet(\mathbb{F}_p[\![G_K(p)]\!])$ is Koszul (cf. [Wei15b, Question 1],

From Observation 1.1 and Theorem 1.2 we obtain a positive answer to Positselski's and Weigel's conjectures in the case of one-relator maximal pro-p Galois groups.

Theorem 1.3. *Let K be a field satisfying the assumptions of Theorem 1.2. Then the algebras $H^\bullet(G_K(p),\mathbb{F}_p)$ and $\mathrm{gr}_\bullet(\mathbb{F}_p[\![G_K(p)]\!])$ are Koszul.*

Finally, we observe that there are finitely generated free-by-Demushkin pro-p groups for which the results of the present paper do hold, but which are not realizable as maximal pro-p Galois groups of fields (cf. Example 5.3). We ask whether one may find further conditions that the defining relation of such pro-p groups should satisfy to make them Galois — or obstructions which prevent them to be maximal pro-p Galois groups.

2. Quadratic algebras and Koszul algebras

2.1. **Graded algebras and Koszul algebras.** Henceforth every graded algebra $A_\bullet$ over a field $\Bbbk$ is assumed to be unitary associative and non-negatively graded of finite-type, i.e., $A_0=\Bbbk$, $A_n=0$ for $n<0$ and $\dim(A_n)<\infty$ for $n\geq 1$.

Let consider $\Bbbk$ as a $A_\bullet$-module via the augmentation epimorphism $\varepsilon\colon A_\bullet\to\Bbbk$. Then one may define the extension spaces $\mathrm{Ext}^n_{A_\bullet}(\Bbbk,\Bbbk)$, $n\geq 0$, as derived functor of the functor $\mathrm{Hom}_{A_\bullet}(\Bbbk,\Bbbk)$ in the usual sense. The cup product (or Yoneda product) induces a graded algebra structure on $H^\bullet(A_\bullet)=\bigoplus_{n\geq 0}\mathrm{Ext}^n_{A_\bullet}(\Bbbk,\Bbbk)$ (cf. [Pri70, § 1] and [PP05, § 1.1]).

The grading of $A_\bullet$ induces an additional internal grading, so that $H^\bullet(A_\bullet)$ may be viewed as a bigraded $\Bbbk$-algebra

$$H^\bullet(A_\bullet)=\bigoplus_{i,j\geq 0}\mathrm{Ext}^{i,j}_{A_\bullet}(\Bbbk,\Bbbk),$$

where $\mathrm{Ext}^{i,j}_{A_\bullet}(\Bbbk,\Bbbk)=0$ for $i>j$ (cf. [PP05, Prop. 1.3.1]). A graded $\Bbbk$-algebra is called a *Koszul* algebra if its extension spaces are cocentrated on the diagonal, i.e., if $\mathrm{Ext}^{i,j}_{A_\bullet}(\Bbbk,\Bbbk)=0$ for $i\neq j$, so that in fact $H^\bullet(A_\bullet)=\bigoplus_{i\geq 0}\mathrm{Ext}^{i,i}_{A_\bullet}(\Bbbk,\Bbbk)$. For further details on the cohomology of graded algebras we refer to [Pri70, § 1-2], [PP05, § 1.1-1.5] and [Pos05, § 2].

2.2. **Quadratic algebras.** For a vector space V over $\Bbbk$ (of finite dimension), let $T_\bullet(V)$ denote the graded tensor $\Bbbk$-algebra generated by V. I.e.,

$$T_\bullet(V) = \bigoplus_{n\geq 0} V^{\otimes n}, \qquad \text{with } V^{\otimes 0} = \Bbbk,$$

endowed with the multiplication induced by the tensor product. Moreover, let $V^* = \mathrm{Hom}_\Bbbk(V, \Bbbk)$ be the $\Bbbk$-dual space of V. Since V has finite dimension, one may identify $(V \otimes V)^* = V^* \otimes V^*$.

Definition 2.1. (i) A graded $\Bbbk$-algebra $A_\bullet$ is said to be *quadratic* if $A_\bullet$ is a quotient $T_\bullet(A_1)/\langle\Omega\rangle$, where $\langle\Omega\rangle$ is the two-sided ideal of $T_\bullet(A_1)$ generated by a subset $\Omega \subseteq A_1 \otimes A_1$.

(ii) For a graded $\Bbbk$-algebra $A_\bullet$, the *quadratic part* of $A_\bullet$, denoted by $\mathrm{q}A_\bullet$, is the quadratic $\Bbbk$-algebra such that $\mathrm{q}A_1 = A_1$ and $\mathrm{q}A_2$ embeds in A_2.

(iii) For a quadratic $\Bbbk$-algebra $A_\bullet = T_\bullet(A_1)/\langle\Omega\rangle$, let $\Omega^\perp \subseteq (A_1 \otimes A_1)^*$ denote the orthogonal space of Ω, i.e.,

$$\Omega^\perp = \{\alpha \in (A_1 \otimes A_1)^* \mid \alpha(\omega) = 0 \text{ for all } \omega \in \Omega\}.$$

Thus, we may consider $\Omega^\perp$ as a subspace of $A_1^* \otimes A_1^*$. The *quadratic dual* of $A_\bullet$, denoted by $A_\bullet^!$, is the quadratic $\Bbbk$-algebra obtained via the quotient $T_\bullet(A_1^*)/\langle\Omega^\perp\rangle$.

In particular, for a quadratic $\Bbbk$-algebra $A_\bullet$ one has $(A_\bullet^!)^! = A_\bullet$. Moreover, one has the following (cf. [PP05, Prop. 1.3.1 and Definition 2.1]).

Proposition 2.2. *let $A_\bullet$ be a graded $\Bbbk$-algebra.*

(i) *The diagonal of $H^\bullet(A_\bullet)$ is isomorphic to the quadratic dual of $\mathrm{q}A_\bullet$, i.e., one has an isomorphism $\bigoplus_{i\geq 0} \mathrm{Ext}^{i,i}_{A_\bullet}(\Bbbk, \Bbbk) \simeq (\mathrm{q}A_\bullet)^!$.*

(ii) *If $A_\bullet$ is Koszul, then it is quadratic.*

The above proposition implies that a quadratic $\Bbbk$-algebra $A_\bullet$ is Koszul if, and only if, one has $H^\bullet(A_\bullet) \simeq A_\bullet^!$. Moreover, a quadratic $\Bbbk$-algebra $A_\bullet$ is Koszul if, and only if, the quadratic dual $A_\bullet^!$ is Koszul (cf. [LV12, Prop. 3.4.8]). For further details on Koszul algebras we refer to [PP05, Ch. 2].

One has the following examples of Koszul $\Bbbk$-algebras (cf. [LV12, Ex.s 3.2.5 and Ex.s 3.4.12]).

Example 2.3. Let V be a vector space of finite dimension over $\Bbbk$.

(a) The tensor $\Bbbk$-algebra $T_\bullet(V)$ and the trivial quadratic $\Bbbk$-algebra $\Bbbk \oplus V$ (with V the part of degree 1) are Koszul algebras, and $T_\bullet(V)^! = \mathbb{F}_p \oplus V^*$ and conversely.

(b) The symmetric algebra $S_\bullet(V)$ and the exterior algebra $\Lambda_\bullet(V)$ are Koszul algebras, and $S_\bullet(V)^! = \Lambda_\bullet(V^*)$ and conversely.

Given two quadratic $\Bbbk$-algebras $A_\bullet$ and $B_\bullet$, the direct product and the free product of $A_\bullet$ and $B_\bullet$ are defined in the following way.

(a) The direct product of $A_\bullet$ and $B_\bullet$ is the quadratic $\mathbb{F}_p$-algebra $C_\bullet = A_\bullet \sqcap B_\bullet$ with $C_n = A_n \oplus B_n$ for every $n \geq 1$.

(b) The free product of $A_\bullet$ and $B_\bullet$ is the quadratic $\mathbb{F}_p$-algebra $C_\bullet = A_\bullet \sqcup B_\bullet$ with $C_1 = A_1 \oplus B_1$ and $C_2 = A_2 \oplus B_2 \oplus (A_1 \otimes B_1) \oplus (B_1 \otimes A_1)$.

For such algebras one has the following (cf. [PP05, § 3.1]).

Proposition 2.4. *Let $A_\bullet$ and $B_\bullet$ be quadratic $\Bbbk$-algebras. If both $A_\bullet$ and $B_\bullet$ are Koszul algebras, then also the direct product $A_\bullet \sqcap B_\bullet$ and the free product $A_\bullet \sqcup B_\bullet$ are Koszul algebras. Moreover, one has*

$$(A_\bullet \sqcap B_\bullet)^! = A_\bullet^! \sqcup B_\bullet^! \qquad \text{and} \qquad (A_\bullet \sqcup B_\bullet)^! = A_\bullet^! \sqcap B_\bullet^!. \tag{2.1}$$

2.3. **Demushkin algebras.** For a vector space V over $\Bbbk$ of finite dimension d, let $X = \{X_1, \dots, X_d\}$ be a set of (non-commutative) indeterminates. The free associative algebra $\Bbbk\langle X\rangle$ — i.e., the $\Bbbk$-algebra of polynomials on the non commutative indeterminates X — comes endowed with the grading induced by the subspaces of homogeneous polynomials. We may identify X with a fixed basis of V, and such identification induces an isomorphism of quadratic $\Bbbk$-algebras $T_\bullet(V) \simeq \Bbbk\langle X\rangle$.

Lemma 2.5. *Let $f \in \Bbbk\langle X\rangle$ be a homogeneous polynomial of degree 2 such that $f \neq X_1^{2\delta_1} + \ldots + X_d^{2\delta_d}$, with $\delta_i \in \{0,1\}$ for every $i = 1, \dots, d$ — i.e., f contains a monomial $\alpha X_i X_j$ with $\alpha \in \Bbbk \smallsetminus \{0\}$ and $i \neq j$. Then the quadratic $\Bbbk$-algebra $\Bbbk\langle X\rangle/\langle f\rangle$ is Koszul.*

Proof. Up to renumbering of X we may assume that f contains the monomial $X_d X_{d-1}$. For every $n \geq 1$ let the monomial basis $\mathcal{S}_n = \{X_{i_1} \cdots X_{i_n}, 1 \leq i_1, \dots, i_n \leq d\}$ be endowed with the lexicographic ordering $\prec$. Then $(\bigcup_n \mathcal{S}_n, \prec)$ is a totally ordered set. In particular, we may write the relation $f = 0$ as

$$X_d X_{d-1} = \text{smaller terms w.r.t. } \prec .$$

In a quadratic $\Bbbk$-algebra $A_\bullet = \Bbbk\langle X\rangle/\langle\Omega\rangle$, with Ω a set of homogeneous polynomials of degree 2, a monomial $h = X_{i_1}X_{i_2}X_{i_3} \in \mathcal{S}_3$ is said to be *critical* if both $X_{i_1}X_{i_2}$ and $X_{i_2}X_{i_3}$ are leading terms (with respect to $\prec$) of some relations in Ω (c.f. [LV12, § 4.1]) — i.e., the monomial h may be rewritten in $A_\bullet$ using lower monomials with respect to $\prec$ in at least two ways, which may eventually lead to different results.

The $\Bbbk$-algebra $\Bbbk\langle X\rangle/\langle f\rangle$ has no critical monomials, as one can not have $X_{i_1}X_{i_2} = X_{i_2}X_{i_3} = X_d X_{d-1}$ for any $i_1, i_2, i_3 \in \{1, \dots, d\}$, and no monomials can be rewritten using lower monomials. By [LV12, Thm. 4.1.1], quadratic $\Bbbk$-algebras without critical monomials are Koszul, and so is $\Bbbk\langle X\rangle/\langle f\rangle$. □

Fix a prime number p. Henceforth we shall concentrate on graded algebras over the finite field $\mathbb{F}_p$. A finitely generated pro-p group G is called a *Demushkin group* if $H^2(G, \mathbb{F}_p) \simeq \mathbb{F}_p$ and the cup product induces a non-degenerate $\mathbb{F}_p$-pairing

$$H^1(G, \mathbb{F}_p) \times H^1(G, \mathbb{F}_p) \longrightarrow H^2(G, \mathbb{F}_p) \simeq \mathbb{F}_p.$$

Demushkin groups are one-relator pro-p groups, and their structure was fully described by S.P. Demushkin, J.-P. Serre and J.P. Labute (cf. [NSW08, § III.9]). Moreover, they appear in Galois theory as the maximal pro-p Galois groups of certain p-adic local fields (cf. [NSW08, Thm. 7.5.11]).

Definition 2.6. A *Demushkin $\mathbb{F}_p$-algebra* $D_\bullet$ is a quadratic $\mathbb{F}_p$-algebra such that its diagonal $\mathbb{F}_p$-cohomology is isomorphic to the $\mathbb{F}_p$-cohomology ring of a Demushkin group. I.e., $\operatorname{Ext}_{D_\bullet}^{1,1} \simeq D_1^*$, $\operatorname{Ext}_{D_\bullet}^{2,2}(\mathbb{F}_p, \mathbb{F}_p) \simeq \mathbb{F}_p$ and the cup product induces a non-degenerate $\mathbb{F}_p$-pairing

$$\operatorname{Ext}_{D_\bullet}^{1,1}(\mathbb{F}_p, \mathbb{F}_p) \times \operatorname{Ext}_{D_\bullet}^{1,1}(\mathbb{F}_p, \mathbb{F}_p) \longrightarrow \mathbb{F}_p.$$

By Proposition 2.2 the diagonal cohomology of a Demushkin $\mathbb{F}_p$-algebra is a quadratic $\mathbb{F}_p$-algebra $A_\bullet$ with $\dim(A_1) = \dim(D_1) = d$, with d even if p is odd. In particular, there is a basis $\{a_1 \ldots, a_d\}$ of A_1 such that

$$a_1a_2 = -a_2a_1 = a_3a_4 = -a_4a_3 = \ldots = a_{d-1}a_d = -a_da_{d-1} \neq 0 \tag{2.2}$$

(and possibly $a_1^2 = a_1a_2$ if $p = 2$), and $a_ia_j = 0$ in any other case, if d is even; and

$$a_1^2 = a_2a_3 = a_3a_2 = a_4a_5 = a_5a_4 = \ldots = a_{d-1}a_d = a_da_{d-1} \neq 0 \tag{2.3}$$

and $a_ia_j = 0$ in any other case, if d is even (and thus necessarily $p = 2$). Demushkin $\mathbb{F}_p$-algebras are described by the following.

Proposition 2.7. *For a Demushkin algebra $D_\bullet$ set $d = \dim(D_1)$. One may present $D_\bullet$ as a quotient $D_\bullet = \mathbb{F}_p\langle X\rangle/\langle f\rangle$, with $X = \{X_1, \ldots, X_d\}$ non-commutative indeterminates and f a homogeneous polynomial of degree two, such that:*

(a) *if p is odd then d is necessarily even and*
$$f = [X_1, X_2] + [X_3, X_4] + \ldots + [X_{d-1}, X_d];$$
(b) *if $p = 2$ and d is even then either f is as above or*
$$f = X_1^2 + [X_1, X_2] + [X_3, X_4] + \ldots + [X_{d-1}, X_d];$$
(c) *if $p = 2$ and d is even then*
$$f = X_1^2 + [X_2, X_3] + [X_4, X_5] + \ldots + [X_{d-1}, X_d].$$

(Here $[X_i, X_j]$ denotes the commutator polynomial $X_iX_j - X_jX_i$.)

Lemma 2.8. *Let $D_\bullet$ be a quadratic $\mathbb{F}_p$-algebra described in Proposition 2.7. Then the quadratic dual $D_\bullet^!$ is a quadratic $\mathbb{F}_p$-algebra $A_\bullet$ described by* (2.2) *or* (2.3).

Proof. Fix a basis $\{a_1 \ldots, a_d\}$ of $D_1^! = D_1^*$ which is dual to X — i.e., $a_i(X_j) = \delta_{ij}$. Let $\Omega \subseteq D_1 \otimes D_1$ be the subspace generated by the homogeneous polynomial f. Then

$$\dim(\Omega) = \dim(A_1 \otimes A_1) - \dim(\Omega^\perp) = \dim(A_2) = 1. \tag{2.4}$$

The only (monic) monomials of $D_1^! \otimes D_1^!$ whose evaluation on f is not trivial are $a_{2h-1}a_{2h}$ and $a_{2h}a_{2h-1}$, with $1 \leq h \leq d/2$, if d is even (and possibly also a_1^2 if the monomial X_1^2 appears in f), or a_1^2, $a_{2h}a_{2h+1}$ and $a_{2h+1}a_{2h}$, with $1 \leq h \leq (d-1)/2$ if d is odd. Therefore, (2.4) implies the relations in $D_2^!$

$$a_{2h-1}a_{2h} = -a_{2h}a_{2h-1} \neq 0, \qquad 1 \leq h \leq \frac{d}{2},$$

and $a_ia_j = 0$ else, if d is even (and possibly a_1^2 equal to the above monomials), and

$$a_1^2 = a_{2h}a_{2h+1} = a_{2h+1}a_{2h} \neq 0, \qquad 1 \leq h \leq \frac{d-1}{2},$$

and $a_ia_j = 0$ else, if d is odd.

These are the same relations as in (2.2) and (2.3), and the claim follows. □

Proof of Proposition 2.7. Let $D_\bullet$ be a Demushkin $\mathbb{F}_p$-algebra, and let $A_\bullet$ be its diagonal cohomology. Then by Proposition 2.2 one has $D_\bullet = (D_\bullet^!)^! = A_\bullet^!$, and the claim follows by Lemma 2.8. □

Proposition 2.9. *A Demushkin $\mathbb{F}_p$-algebra $D_\bullet$ is Koszul.*

Proof. Assume first that $p = 2$ and $\dim(D_1) = 1$. Then $D_\bullet = \mathbb{F}_p[X_1]/\langle X_1^2\rangle$, i.e., $D_\bullet$ is a trivial quadratic $\mathbb{F}_2$-algebra, and such algebra is Koszul by Example 2.3. If $\dim(D_1) > 1$ then the claim follows by Proposition 2.7 and Lemma 2.5. □

3. Pro-p groups and graded algebras

3.1. **The Zassenhaus filtration.** Henceforth, subgroups of pro-p groups are assumed to be closed (in the pro-p topology), and every generating set is to be intended in the topological sense. In particular, given two (closed) subgroups C_1 and C_2 of a pro-p group G, the subgroup $[C_1, C_2]$ is the (closed) subgroup of G generated by the commutators

$$[x, y] = (y^{-1})^x \cdot y = x^{-1}y^{-1}xy, \qquad x \in C_1, y \in C_2.$$

Moreover, for a closed subgroup C of G and $n \geq 1$, C^n denotes the (closed) subgroup of G generated by the elements g^n, with $g \in C$.

For a pro-p group G let $\mathbb{F}_p[\![G]\!]$ be the *complete* group algebra. I.e.,

$$\mathbb{F}_p[\![G]\!] = \varprojlim_{U \lhd_o G} \mathbb{F}_p[G/U],$$

with U running through the set of open normal subgroups of G. The augmentation ideal $I \subseteq \mathbb{F}_p[\![G]\!]$ is the kernel of the augmentation map $\mathbb{F}_p[\![G]\!] \to \mathbb{F}_p$, given by $g \mapsto 1$ for every $g \in G$. Thus, one defines the *Zassenhaus filtration* of G as follows.

Definition 3.1. The Zassenhaus filtration $G_{(n)}$, $n \geq 1$, is the filtration of normal subgroups of G defined by $G_{(n)} = \{g \in G \mid g - 1 \in I^n\}$ for every $n \geq 1$.

The Zassenhaus filtration of a pro-p group G is the fastest filtration starting at G with the following property: for every $x \in G_{(n)}$ and $y \in G_m$, $n, m \geq 1$, one has

$$[x, y] \in G_{(n+m)} \qquad \text{and} \qquad x^p \in G_{(np)} \tag{3.1}$$

In particular, one has $G_{(1)} = G$, $G_{(2)} = G^p[G, G]$ — i.e., $G_{(2)}$ is the Frattini subgroup of G —, and

$$G_{(3)} = \begin{cases} G^p[[G, G], G] & \text{if } p \neq 2 \\ G^4[G, G]^2[[G, G], G] & \text{if } p = 2 \end{cases} \tag{3.2}$$

(cf. [DdSMS99, § 11.1]).

Remark 3.2. Let F be a finitely generated pro-p group, and let $\{x_1, \ldots, x_d\}$ be a minimal set of generators for F. Moreover, let $\mathbb{F}_p\langle\langle X\rangle\rangle$ be the $\mathbb{F}_p$-algebra of formal power series on the set of indeterminates $X = \{X_1, \ldots, X_d\}$. Then one has an isomorphism of (topological) $\mathbb{F}_p$-algebras $\mathbb{F}_p[\![F]\!] \simeq \mathbb{F}_p\langle\langle X\rangle\rangle$, called the *Magnus morphism*, given by $x_i \mapsto 1 + X_i$ for every $i = 1, \ldots, d$. The Magnus morphism maps the augmentation ideal of $\mathbb{F}_p[\![F]\!]$ onto the two-sided ideal

$$I(X) = \langle X_1, \ldots, X_d\rangle = \bigoplus_{i=1}^{d} \mathbb{F}_p\langle\langle X\rangle\rangle \cdot X_i.$$

Thus, we may identify the graded algebra $\mathrm{gr}_\bullet(\mathbb{F}_p\langle\langle X\rangle\rangle) = \bigoplus_{n\geq 0} I(X)^n/I(X)^{n+1}$, with $I(X)^0 = \mathbb{F}_p\langle\langle X\rangle\rangle$, with the free polynomial algebra $\mathbb{F}_p\langle X\rangle$, endowed with the grading induced by the subspaces of homogeneous polynomials. Also, via this identification one has $F_{(n)} = \{g \in F \mid g - 1 \in I(X)^n\}$ for every $n \geq 1$.

3.2. **Pro-p groups and restricted Lie algebras.** Given an associative $\mathbb{F}_p$-algebra R, let R_L denote the Lie $\mathbb{F}_p$-algebra endowed with the Lie brackets $(a,b) = ab - ba$ for every $a, b \in R$. One has the following definition (cf. [DdSMS99, § 12.1]).

Definition 3.3. (i) A *restricted* Lie $\mathbb{F}_p$-algebra L over $\mathbb{F}_p$ is a Lie $\mathbb{F}_p$-algebra equipped with a p-power map $[p]\colon L \longrightarrow L$ which is compatible with the Lie brackets. I.e., there exists an associative $\mathbb{F}_p$-algebra R and a monomorphism of Lie algebras $\theta\colon L \to R_L$ such that $\theta(v^{[p]}) = \theta(v)^p$ for all $v \in L$.

(ii) The algebra R, endowed with the monomorphism $\theta\colon L \to R_L$, is called the *universal restricted envelope* of L if for any morphism of restricted Lie algebras $\varphi\colon L \to B_L$, with B an associative $\mathbb{F}_p$-algebra, there exists a unique homomorphism of associative $\mathbb{F}_p$-algebras $\tilde{\varphi}\colon R \to B$ such that $\varphi = \tilde{\varphi} \circ \theta$.

We shall denote the universal restricted envelope of a restricted Lie $\mathbb{F}_p$-algebra L by $\mathcal{U}(L)$. A restricted ideal $\mathfrak{r}$ of a restricted Lie $\mathbb{F}_p$-algebra is an ideal in the classic sense with the further condition that $v^{[p]} \in \mathfrak{r}$ for every $v \in \mathfrak{r}$. Then one has the following (cf. [Gär15, Prop. 2.1])

Proposition 3.4. *Let L be a restricted Lie algebra over $\mathbb{F}_p$ and let $\mathfrak{r} \subseteq L$ be a restricted ideal. Let $\mathcal{R}$ be the left ideal of $\mathcal{U}(L)$ generated by the image of $\mathfrak{r}$ via the embedding $\theta\colon L \to \mathcal{U}(L)$. Then $\mathcal{R}$ is a two-sided ideal and the epimorphism $L \to L/\mathfrak{r}$ induces a short exact sequence*

$$0 \longrightarrow \mathcal{R} \longrightarrow \mathcal{U}(L) \longrightarrow \mathcal{U}(L/\mathfrak{r}) \longrightarrow 0 \tag{3.3}$$

For a pro-p group G let $L(G)$ be the graded object

$$L(G) = \bigoplus_{n \geq 1} G_{(n)}/G_{(n+1)}.$$

By (3.1) every quotient $G_{(n)}/G_{(n+1)}$ is a $\mathbb{F}_p$-vector space, and the commutators and the p-power induce the structure of graded restricted Lie algebra on $L(G)$. Moreover, the universal restricted envelope of $L(G)$ is the graded $\mathbb{F}_p$-algebra $\mathrm{gr}_\bullet(\mathbb{F}_p[\![G]\!])$ (cf. [DdSMS99, Thm. 12.8]). If $g \in G_{(n)} \smallsetminus G_{(n+1)}$, $n \geq 1$, we shall call the image of g in $G_{(n)}/G_{(n+1)}$ the *initial form* of g in $L(G)$.

Remark 3.5. If F is a finitely generated free pro-p group, with minimal generating set $\{x_1, \ldots, x_d\}$, we may identify the universal restricted envelope $\mathcal{U}(L(F))$ — and therefore the graded group algebra $\mathrm{gr}_\bullet(\mathbb{F}_p[\![F]\!])$ — with the free polynomial algebra $\mathbb{F}_p\langle X\rangle$, with $X = \{X_1, \ldots, X_d\}$. In particular, the restricted Lie $\mathbb{F}_p$-algebra $L(F)$ is the free restricted Lie $\mathbb{F}_p$-algebra on X (cf. [Gär15, Remark 2.3]).

Let $G = F/R$ be a finitely generated pro-p group, with R a normal subgroup of F. Then the restricted Lie $\mathbb{F}_p$-algebra $L(G)$ is a quotient $L(F)/\mathfrak{r}$, with $\mathfrak{r}$ a restricted ideal of $L(F)$, and thus by Proposition 3.4 the graded $\mathbb{F}_p$-algebra $\mathrm{gr}_\bullet(\mathbb{F}_p[\![G]\!])$ is isomorphic to a quotient $\mathbb{F}_p\langle X\rangle/\mathcal{R}$, with $\mathcal{R}$ the two-sided ideal of $\mathbb{F}_p\langle X\rangle$ generated by $\mathfrak{r}$.

3.3. **Mild pro-p groups.** Moreover, such a pro-p group G is called *mild* (with respect to the Zassenhaus filtration) if there is a minimal set of defining relations $\{r_1, \ldots, r_m\}$ (i.e., a minimal subset of F which generates R as normal subgroup of F) which is a *strongly free* sequence (with respect to the Zassenhaus filtration), see [Gär15, Definition 2.7 and Definition 2.11]. Mild pro-p groups were introduced by

J. Labute in [Lab06] to study the Galois groups of the extensions of number fields with restricted ramification. Mild pro-p groups have the following properties (cf. [Gär15, Thm. 2.12]).

Proposition 3.6. *Let G be a mild pro-p group, with a strongly free minimal set of defining relations $\{r_1, \dots, r_m\}$. Then*

(i) *one has $H^n(G) = 0$ for all $n > 2$;*
(ii) *one has an isomorphism of graded $\mathbb{F}_p$-algebras*

$$\mathrm{gr}_\bullet(\mathbb{F}_p[\![G]\!]) \simeq \mathbb{F}_p\langle X\rangle/\mathcal{R}, \tag{3.4}$$

where $\mathcal{R}$ denotes the two-sided ideal generated by the initial forms of r_i's.

4. One-relator pro-p groups

4.1. **Cohomology of one-relator pro-p groups.** For a pro-p group G we shall denote the $\mathbb{F}_p$-cohomology groups simply by $H^n(G)$ for every $n \geq 0$. If G is a finitely generated one-relator pro-p group, then one has the minimal presentation

$$1 \longrightarrow R \longrightarrow F \longrightarrow G \longrightarrow 1 \tag{4.1}$$

with $H^1(F) \simeq H^1(G)$ of dimension $d = \dim(H^1(G))$ equal to the minimal number of generators of G, and R generated as normal subgroup of F by an element $r \in F$, which is called the defining relation. In particular, one has $r \in F_{(2)}$, as the presentation is minimal, and $H^2(G) \simeq \mathbb{F}_p$, as $\dim(H^2(G))$ is the minimal number of generators of R as normal subgroup of F (cf. [NSW08, § III.9]).

Proposition 4.1. *Let $G = F/R$ be a finitely generated one-relator pro-p group, with defining relation $r \in F_{(2)} \smallsetminus F_{(3)}$ (if $p = 2$ assume further that $r \notin F_{(2)} \smallsetminus [F, F]$). Then G is mild, and the $\mathbb{F}_p$-cohomology ring $H^\bullet(G)$ is quadratic.*

Proof. By hypothesis, one may find a a minimal generating set $\{x_1, \dots, x_d\} \subseteq F$ such that the commutator $[x_i, x_j]$ appears in r for some $i, j \in \{1, \dots, d\}$, $i \neq j$. Up to renumbering, we may assume $i = d - 1$ and $j = d$.

Identify the graded group algebra $\mathrm{gr}_\bullet(\mathbb{F}_p[\![F]\!])$ with the algebra $\mathbb{F}_p\langle X\rangle$, with $X = \{X_1, \dots, X_d\}$. The initial form ρ of r is a homogeneous polynomial of degree two, with leading monomial $\alpha X_d X_{d-1}$, with $\alpha \in \mathbb{F}_p$. Since the sequence $\{X_n X_{n-1}\}$ is combinatorially free in the sense of [Gär15, Definition 3.1], the sequence $\{\rho\} \subseteq \mathbb{F}_p\langle X\rangle$ is strongly free by [Gär15, Thm. 3.5], and G is a mild. (In the case $p \neq 2$ this follows also from [Gär15, Cor. 5.10].)

Moreover, by [NSW08, Prop. 3.9.13] the one has $\chi_{d-1} \cup \chi_d \neq 0$. Thus, $\chi_{d-1} \cup \chi_d$ generates $H^2(G) \simeq \mathbb{F}_p$, and $H^n(G) = 0$ for $n > 2$ by [Gär15, Thm. 2.12-(i)]. Therefore, $H^\bullet(G)$ is a quadratic $\mathbb{F}_p$-algebra. □

Henceforth, we shall assume that the $\mathbb{F}_p$-cohomology ring $H^\bullet(G)$ is quadratic. The cup product induces a (skew-commutative) $\mathbb{F}_p$-pairing

$$H^1(G) \times H^1(G) \xrightarrow{\cup} \mathbb{F}_p\ . \tag{4.2}$$

(cf. [NSW08, Prop. 3.9.12]). If the pairing (4.2) is perfect, then G is a Demushkin group (cf. [NSW08, Definition 3.9.9]). Otherwise, let $V_2 = H^1(G)^\perp$ be the radical of $H^1(G)$ with respect to the cup product — i.e.,

$$V_2 = H^1(G)^\perp = \{\chi \in H^1(G) \mid \chi \cup \psi = 0 \text{ for all } \psi \in H^1(G)\}.$$

Then we may decompose $H^1(G) = V_1 \oplus V_2$, so that (4.2) induces a perfect pairing $V_1 \times V_1 \to \mathbb{F}_p$.

Let $G^{ab} = G/[G,G]$ be the abelianization of G. Since G is a one-relator pro-p group, one has an isomorphism of abelian pro-p groups

$$G^{ab} \simeq \mathbb{Z}_p^d \qquad \text{or} \qquad G^{ab} \simeq \mathbb{Z}_p/q\mathbb{Z}_p \times \mathbb{Z}_p^{d-1},$$

with $d = \dim(H^1(G))$ and q a power of p. In the former case we shall say that $q = 0$. Now one has two cases:

(i) p is odd, or $p = 2$ and $q \neq 2$;
(ii) $p = q = 2$.

4.2. **First case: $q \neq 2$.** Set $n = \dim(V_1)$ and $m = \dim(V_2)$, and let (4.1) be a minimal presentation of G. By [NSW08, Prop. 3.9.13] we may choose a minimal generating set $\mathcal{S} = \{x_1, \dots, x_n, y_1, \dots, y_m\}$ of F such that

$$r \equiv \prod_{i=1}^{n} x_i^{a_i p} \cdot \prod_{i=1}^{m} y_i^{b_i p} \cdot \prod_{1 \leq i < j \leq n} [x_i, x_j]^{a_{ij}} \mod F_{(2)}^p [F_{(2)}, F], \tag{4.3}$$

with $0 \leq a_i, b_i, a_{ij} \leq p-1$ for every i, j (such that $a_i p, b_i p \geq q$ for every i). By (3.2), from (4.3) one obtains

$$r \equiv \prod_{1 \leq i < j \leq n} [x_i, x_j]^{a_{ij}} \mod F_{(3)}. \tag{4.4}$$

Since $q \neq 2$, the pairing (4.2) is alternating (see also [QW, § 3.2]), and thus n is even and V_1 decomposes into a direct sum of hyperbolic planes. Therefore, one may find a basis $\mathcal{B}_1 = \{\chi_1, \dots, \chi_n\}$ of V_1 with χ_1 dual to x_1, which completes to a bases $\mathcal{B}$ of $H^1(G)$ such that

$$\chi_1 \cup \chi_2 = \chi_3 \cup \chi_4 = \ldots = \chi_{n-1} \cup \chi_n = 1 \qquad \text{in } H^2(G) \simeq \mathbb{F}_p, \tag{4.5}$$

and $\chi \cup \psi = 0$ in any other case for $\chi, \psi \in \mathcal{B}$ (cf. [NSW08, Prop. 3.9.16]). Therefore, one has the following.

Proposition 4.2. *The $\mathbb{F}_p$-cohomology ring of G splits as direct product*

$$H^\bullet(G) = A_\bullet \sqcap B_\bullet, \tag{4.6}$$

with $A_1 = V_1$ and $A_2 \simeq H^2(G)$, and $B_\bullet$ the trivial quadratic $\mathbb{F}_p$-algebra generated by V_2.

Proof. By definition one has $H^1(G) = A_1 \oplus B_1$. Also, $B_2 = A_n = B_n = 0$ for every $n \geq 3$, so that $H^n(G) = A_n \oplus B_n$ for $n \geq 2$. In particular, by (4.5) the cup product induces a perfect pairing $A_1 \times A_1 \to \mathbb{F}_p$. □

Without loss of generality, we may choose the minimal generating set $\mathcal{S}$ to be the dual of to $\mathcal{B}$ Then by [NSW08, Prop. 3.9.13] equation (4.4) becomes

$$r \equiv [x_1, x_2][x_3, x_4] \cdots [x_{n-1}, x_n] \mod F_{(3)}. \tag{4.7}$$

Therefore, under the identification $\mathcal{U}(L(F)) = \mathbb{F}_p\langle X \rangle$, with $X = \{X_1, \dots, X_d\}$, the initial form of r in $F_{(2)}/F_{(3)}$ is

$$\rho = [X_1, X_2] + [X_3, X_4] + \cdots + [X_{n-1}, X_n] \in \mathbb{F}_p\langle X \rangle. \tag{4.8}$$

By Proposition 4.1, G is mild. Therefore, the restricted Lie algebra $L(G)$ is the quotient $L(X)/\mathfrak{r}$, where $\mathfrak{r}$ is the restricted ideal of $L(X)$ generated by ρ, and the

two-sided ideal $\mathcal{R}$ of $\mathcal{U}(L(X))$ generated by $\mathfrak{r}$ is the two-sided ideal generated by ρ. Thus, Proposition 3.4 implies the following.

Proposition 4.3. *In this case the graded algebra* $\mathrm{gr}_\bullet(\mathbb{F}_p[\![G]\!])$ *splits as free product:*

$$\mathrm{gr}_\bullet(\mathbb{F}_p[\![G]\!]) = D_\bullet \sqcup \mathbb{F}_p\langle X'\rangle, \tag{4.9}$$

with $D_\bullet$ *the Demushkin quadratic* $\mathbb{F}_p$*-algebra with* $\dim(D_1) = n$ *and defining relation* ρ *as in* (4.8)*, and* $\mathbb{F}_p\langle X'\rangle$ *the free polynomial algebra on* $X' = \{X_{n+1}, \dots, X_d\}$.

4.3. **Second case:** $q = 2$**.** Set $n = \dim(V_1)$ and $m = \dim(V_2)$, and let (4.1) be a minimal presentation of G. Assume $n \geq 2$. By [NSW08, Prop. 3.9.13] and by (3.2) we may choose a minimal generating set $\mathcal{S} = \{x_1, \dots, x_n, y_1, \dots, y_m\}$ of F such that $\bar{x}_1^2 \equiv 1 \bmod [G, G]$ (where $\bar{x}_1$ denotes the image of x_1 in G), and

$$r \equiv x_1^2 \prod_{1 \leq i < j \leq n} [x_i, x_j]^{a_{ij}} \mod F_{(3)}, \tag{4.10}$$

with $a_{ij} \in \{0, 1\}$ for every i, j.

Let $\mathcal{B}_1 = \{\chi_1, \dots, \chi_n\}$ be a basis of V_1 with χ_1 dual to x_1. Then again by [NSW08, Prop. 3.9.13] one has $\chi_1 \cup \chi_1 = 1$, and n can be both odd or even. Thus, by [Lab67, Prop. 4] we may choose the basis $\mathcal{B}_1$ and complete it to a basis $\mathcal{B}$ of $H^1(G)$ such that

$$\begin{aligned} \chi_1 \cup \chi_2 = \chi_3 \cup \chi_4 = \ldots = \chi_{n-1} \cup \chi_n = 1, &\qquad \text{if } 2 \mid n, \\ \chi_2 \cup \chi_3 = \chi_4 \cup \chi_5 = \ldots = \chi_{n-1} \cup \chi_n = 1, &\qquad \text{if } 2 \nmid n. \end{aligned}$$

and $\chi \cup \psi = 0$ in any other case for $\chi, \psi \in \mathcal{B}$.

Proposition 4.4. *The* $\mathbb{F}_p$*-cohomology ring of* G *splits as direct product*

$$H^\bullet(G) = A_\bullet \sqcap B_\bullet, \tag{4.11}$$

with $A_1 = V_1$ *and* $A_2 \simeq H^2(G)$*, and* $B_\bullet$ *the trivial quadratic* $\mathbb{F}_p$*-algebra generated by* V_2.

Proof. By definition one has $H^1(G) = A_1 \oplus B_1$. Also, $B_2 = A_n = B_n = 0$ for every $n \geq 3$, so that $H^n(G) = A_n \oplus B_n$ for $n \geq 2$. In particular, by (4.5) the cup product induces a perfect pairing $A_1 \times A_1 \to \mathbb{F}_p$. □

Let the minimal generating set $\mathcal{S}$ be the dual to $\mathcal{B}$. By [Lab67, p. 116] from (4.10) one obtains

$$r \equiv \begin{cases} x_1^2[x_1, x_2][x_3, x_4] \cdots [x_{n-1}, x_n] \mod F_{(3)} & \text{if } 2 \mid n, \\ x_1^2[x_2, x_3][x_4, x_5] \cdots [x_{n-1}, x_n] \mod F_{(3)} & \text{if } 2 \nmid n. \end{cases} \tag{4.12}$$

Therefore, under the identification $\mathcal{U}(L(F)) = \mathbb{F}_2\langle X\rangle$, with $X = \{X_1, \dots, X_d\}$, the initial form of r in $F_{(2)}/F_{(3)}$ is

$$\rho = \begin{cases} X_1^2 + [X_1, X_2] + [X_3, X_4] + \cdots + [X_{n-1}, X_n] & \text{if } 2 \mid n, \\ X_1^2 + [X_2, X_3] + [X_4, X_5] + \ldots + [X_{n-1}, X_n] & \text{if } 2 \nmid n. \end{cases} \tag{4.13}$$

in $\mathbb{F}_p\langle X\rangle$.

By Proposition 4.1, G is mild. Therefore, the restricted Lie algebra $L(G)$ is the quotient $L(X)/\mathfrak{r}$, where $\mathfrak{r}$ is the restricted ideal of $L(X)$ generated by ρ, and the two-sided ideal $\mathcal{R}$ of $\mathcal{U}(L(X))$ generated by $\mathfrak{r}$ is the two-sided ideal generated by ρ. Thus, Proposition 3.4 implies the following.

Proposition 4.5. *In this case the graded algebra* $\mathrm{gr}_\bullet(\mathbb{F}_p[\![G]\!])$ *splits as free product:*

$$\mathrm{gr}_\bullet(\mathbb{F}_2[\![G]\!]) = D_\bullet \sqcup \mathbb{F}_2\langle X'\rangle, \tag{4.14}$$

with $D_\bullet$ *the Demushkin quadratic* $\mathbb{F}_2$*-algebra with* $\dim(D_1) = n$ *and defining relation as in* (4.13)*, and* $\mathbb{F}_2\langle X'\rangle$ *the free polynomial algebra on* $X' = \{X_{n+1}, \dots, X_d\}$*.*

Remark 4.6. If $p = 2$ and $n = 1$ then the initial form of the defining relation r is X_1^2. In particular, explicit computations show that

$$\mathrm{gr}_\bullet(\mathbb{F}_p[\![G]\!]) = \frac{\mathbb{F}_2[X_1]}{\langle X_1^2\rangle} \sqcup \mathbb{F}_2\langle X'\rangle. \tag{4.15}$$

On the other hand, the only cup product which survives in $H^2(G)$ is $\chi_1 \cup \chi_1$, and the $\mathbb{F}_2$-cohomology ring is

$$H^\bullet(G) = \mathbb{F}_2[\chi_1] \sqcap \frac{T_\bullet(V_2)}{\langle V_2 \otimes V_2\rangle}, \tag{4.16}$$

with $\mathbb{F}_2[\chi_1]$ the polynomial ring on χ_1 considered as an indeterminate — i.e., one has $H^k(G) = \mathbb{F}_2\chi_1^k$ for every $k > 1$. This is the only case where $H^k(G)$ is not trivial for $k > 2$ — i.e., the cohomological dimension of G is more than two.

5. Main results

5.1. **Maximal pro-p Galois groups.** Assume that the maximal pro-p Galois group $G = G_K(p)$ of a field K is a finitely generated one-relator pro-p group. Note that G is finitely generated if, and only if, the quotient $K^\times/(K^\times)^p$ of the multiplicative group $K^\times = K \smallsetminus \{0\}$ is finite, as one has the (Pontryagin) duality

$$K^\times/(K^\times)^p = H^1(G, \mathbb{F}_p) = (G/G_{(2)})^*.$$

A pro-p group G is called a *Bloch-Kato pro-p group* if the $\mathbb{F}_p$-cohomology algebra $H^\bullet(C)$ is a quadratic $\mathbb{F}_p$-algebra for every closed subgroup C of G. If K contains a root of unity of order p, the Galois group $G_K(p)$ is a Bloch-Kato pro-p group by the Rost-Voevodsky Theorem (cf. [Qua14, § 2]).

Observation 1.1 follows directly by Proposition 4.2 and Proposition 4.4 (and by Remark 4.6 in the case $p = q = 2$ and $n = 1$).

Proof of Theorem 1.2 and Theorem 1.3. Set $d = \dim(H^1(G_K(p)))$, and let q, and n, m be as in Section 4.

Assume first that $p = q = 2$ and $n = 1$. By (4.15) the graded group algebra $\mathrm{gr}_\bullet(\mathbb{F}_p[\![G_K(p)]\!])$ splits as free product of a trivial quadratic $\mathbb{F}_p$-algebra (on one generator) and a free associative graded $\mathbb{F}_p$-algebra (on m generators). The former is a Demushkin $\mathbb{F}_p$-algebra by Proposition 2.7, and (1.2) holds. Moreover, by (4.16) the $\mathbb{F}_p$-cohomology algebra $H^\bullet(G_K(p))$ decomposes as direct product of the free $\mathbb{F}_p$-algebra on $V_1 \simeq \mathbb{F}_p$ and the trivial quadratic $\mathbb{F}_p$-algebra on V_2. By Example 2.3 such algebras are quadratic dual respectively to the $\mathbb{F}_p$-algebras in (4.15), and thus

$$H^\bullet(G_K(p)) \simeq \mathrm{gr}_\bullet(\mathbb{F}_p[\![G_K(p)]\!])^! \tag{5.1}$$

by Proposition 2.4. This establishes Theorem 1.2 in this case. Finally, by Example 2.3 all the algebras appearing in (4.15) and (4.16) are Koszul, thus both $\mathrm{gr}_\bullet(\mathbb{F}_p[\![G_K(p)]\!])$ and $H^\bullet(G_K(p))$ are Koszul by Proposition 2.4, and this proves Theorem 1.3 in this case.

Assume now that p is odd, or $p = 2$ and $q \neq 2$, so that we are in the first case. The isomorphism (4.9) yields (1.2). Moreover, by Lemma 2.8 the Demushkin $\mathbb{F}_p$-algebra $D_\bullet$ in (4.9) is quadratic dual to the algebra $A_\bullet$ in (4.6), and by Example 2.3 the free $\mathbb{F}_p$-algebra in (4.9) is quadratic dual to the trivial quadratic $\mathbb{F}_p$-algebra $B_\bullet$ in (4.6), so that Proposition 2.4 yields duality (5.1). This establishes Theorem 1.2 in this case. Finally, by Example 2.3 and Proposition 2.9 all such algebras are Koszul, and Proposition 2.4 yields Theorem 1.3 in this case.

At last, assume that $p = q = 2$ and $n > 1$. The isomorphism (4.14) yields (1.2). Moreover, by Lemma 2.8 the Demushkin $\mathbb{F}_p$-algebra $D_\bullet$ in (4.14) is quadratic dual to the algebra $A_\bullet$ in (4.11), and by Example 2.3 the free $\mathbb{F}_p$-algebra in (4.14) is quadratic dual to the trivial quadratic $\mathbb{F}_p$-algebra $B_\bullet$ in (4.11), so that Proposition 2.4 yields duality (5.1). This establishes Theorem 1.2 in this case. Finally, by Example 2.3 and Proposition 2.9 all such algebras are Koszul, and Proposition 2.4 yields Theorem 1.3 in this case. □

Moreover, it is possible to generalize Würfel's theorem to Bloch-Kato pro-p groups as follows.

Theorem 5.1. *Let G be a finitely generated one-relator Bloch-Kato pro-p group. One has the short exact sequence of pro-p groups*

$$1 \longrightarrow N \longrightarrow G \longrightarrow \bar{G} \longrightarrow 1 \tag{5.2}$$

— where N is the normal closure of a free pro-p subgroup $C \subseteq G$, and $\bar{G}$ is a Demushkin group —, which induces the isomorphisms

$$H^\bullet(G) \simeq H^\bullet(\bar{G}) \sqcap H^\bullet(C), \qquad \textit{and}$$
$$\mathrm{gr}_\bullet(\mathbb{F}_p[\![G]\!]) \simeq \mathrm{gr}_\bullet(\mathbb{F}_p[\![\bar{G}]\!]) \sqcup \mathrm{gr}_\bullet(\mathbb{F}_p[\![C]\!]).$$

In particular, this is true when $G = G_K(p)$ is a finitely generated one-relator maximal pro-p Galois group, with K conaining a root of unity of order p.

Proof. Let C be the closed subgroup of G such that the restriction morphism $\mathrm{res}^1_{G,C} \colon H^1(G) \to H^2(C)$ induces an isomorphism $H^1(G)^\perp \simeq H^1(C)$. In particular, one has $\ker(\mathrm{res}^1_{G,C}) = V_1$. Therefore, the commutative diagram

$$\begin{array}{ccc} H^1(G) \times H^1(G) & \xrightarrow{\cup} & H^2(G) \\ \downarrow{\scriptstyle \mathrm{res}^1_{G,C}} \quad \downarrow{\scriptstyle \mathrm{res}^1_{G,C}} & & \downarrow{\scriptstyle \mathrm{res}^2_{G,C}} \\ H^1(C) \times H^1(C) & \xrightarrow{\cup} & H^2(C) \end{array}$$

implies that the lower horizontal arrow is trivial and thus $H^2(C) = 0$, as G is a Bloch-Kato pro-p group. Consequently, C is a free pro-p group (cf. [NSW08, Prop. 3.5.17]).

Let $N \subseteq G$ be the normal closure of C in G, and set $\bar{G} = G/N$. Since $H^1(N)^{\bar{G}} \simeq H^1(C)$, the five terms exact sequence induced by the quotient G/N implies that $H^1(\bar{G}) \simeq V_1$ and that the inflation map $\mathrm{inf}^2_{G,N} \colon H^2(\bar{G}) \to H^2(G)$ is a

monomorphism (cf. [NSW08, Prop. 1.6.7]). Thus, in the commutative diagram

$$\begin{array}{ccc} H^1(\bar{G}) \times H^1(\bar{G}) & \xrightarrow{\cup} & H^2(\bar{G}) \\ \downarrow{\scriptstyle \inf^1_{G,N}} \quad \downarrow{\scriptstyle \inf^1_{G,N}} & & \downarrow{\scriptstyle \inf^2_{G,N}} \\ H^1(G) \times H^1(G) & \xrightarrow{\cup} & H^2(G) \end{array}$$

the upper line is a non-degenerate pairing — in particular, $\bar{G}$ is a one relator pro-p group too. Therefore, $\bar{G}$ is a Demushkin group. □

5.2. **Koszul algebras of elementary type.** In analogy with the class of pro-p groups of elementary type defined by I. Efrat (cf. [Efr98, § 3]), one defines the class of *compact Koszul $\mathbb{F}_p$-algebras of elementary type*, respectively the class of *discrete Koszul $\mathbb{F}_p$-algebras of elementary type*, as the minimal class of quadratic $\mathbb{F}_p$-algebras whose "building blocks" are free associative $\mathbb{F}_p$-algebras and Demushkin $\mathbb{F}_p$-algebras, resp. trivial quadratic $\mathbb{F}_p$-algebras and the $\mathbb{F}_p$-cohomology algebras of Demushkin pro-p groups (i.e., quadratic $\mathbb{F}_p$-algebras $A_\bullet$ subject to (2.2) or (2.3), with $A_n = 0$ for $n \geq 3$), and whose elements can be assembled according to the following rules:

(a) if the Koszul $\mathbb{F}_p$-algebras $A_\bullet$ and $B_\bullet$ are compact of elementary type, then also the free product $A_\bullet \sqcup B_\bullet$ is compact of elementary type,

(b) if the compact Koszul $\mathbb{F}_p$-algebras $A_\bullet$ is of elementary type and $B_\bullet$ is a free algebra with $\dim(B_n) = 1$ for every $n \geq 0$, then also the quadratic algebra

$$A_\bullet \otimes^1 B_\bullet = \frac{A_\bullet \sqcup B_\bullet}{\langle \Omega \rangle}, \quad \text{with } \Omega = \{ab - ba, a \in A_1, b \in B_1\}$$

is compact of elementary type (the operation $\otimes^1$ is called the *commutative tensor product* of quadratic algebras, cf. [PP05, § 3.1]),

for the class of compact Koszul $\mathbb{F}_p$-algebras of elementary type; and

(a') if the Koszul $\mathbb{F}_p$-algebras $A_\bullet$ and $B_\bullet$ are discrete of elementary type, then also the direct product $A_\bullet \sqcap B_\bullet$ is discrete of elementary type,

(b') if the Koszul $\mathbb{F}_p$-algebras $A_\bullet$ is discrete of elementary type and $B_\bullet = \mathbb{F}_p \oplus B_1$ is a trivial quadratic $\mathbb{F}_p$-algebra with $\dim(B_1) = 1$, then also the quadratic algebra

$$A_\bullet \otimes^{-1} B_\bullet = \frac{A_\bullet \sqcup B_\bullet}{\langle \Omega \rangle}, \quad \text{with } \Omega = \{ab + ba, a \in A_1, b \in B_1\}$$

is discrete of elementary type (the operation $\otimes^{-1}$ is called the *skew-commutative tensor product* of quadratic algebras, cf. [PP05, § 3.1]),

for the class of discrete Koszul $\mathbb{F}_p$-algebras of elementary type (cf. [MPQT, § 4.7]).

In [Efr95], Efrat conjectures that if K contains a root of unity of order p and the maximal pro-p Galois group $G_K(p)$ is finitely generated, then $G_K(p)$ is a pro-p group of elementary type. The results of Section 4 — together with Theorem 1.2 — yield the following.

Corollary 5.2. *If G is a finitely generated one-relator pro-p group with a quadratic $\mathbb{F}_p$-cohomology algebra $H^\bullet(G)$, then the graded group algebra $\mathrm{gr}_\bullet(\mathbb{F}_p[\![G]\!])$ is a compact Koszul $\mathbb{F}_p$-algebra of elementary type, and $H^\bullet(G)$ is a discrete Koszul $\mathbb{F}_p$-algebra of elementary type. In particular, this is true for $G = G_K(p)$, with K a field contining a root of unity of order p.*

In [MPQT, Question 2] it is conjectured that the graded group $\mathbb{F}_p$-algebra of $G_K(p)$, resp. the $\mathbb{F}_p$-cohomology algebra $H^\bullet(G_K(p))$, is a compact Koszul $\mathbb{F}_p$-algebra of elementary type, resp. is a discrete Koszul $\mathbb{F}_p$-algebra of elementary type, for every field K containing a root of unity of order p with $G_K(p)$ finitely generated. Thus, Corollary 5.2 provides a positive answer to this conjecture in the case of one-relator maximal pro-p Galois groups.

At the end of [Wür85], Würfel asked whether every one-relator absolute Galois pro-p group — subject to some further cohomological restrictions, as such paper deals with the case where the base field K contains all roots of unity of order a p-power — decomposes as a free pro-p product $\bar{G} *_{\hat{p}} F$ of a Demushkin group $\bar{G}$ with a finitely generated free pro-p group F. In fact, for $G = \bar{G} *_{\hat{p}} F$ one has

$$H^\bullet(G) \simeq H^\bullet(\bar{G}) \sqcap H^\bullet(F) \qquad \text{and}$$
$$\mathrm{gr}_\bullet(\mathbb{F}_p[\![G]\!]) \simeq \mathrm{gr}_\bullet(\mathbb{F}_p[\![\bar{G}]\!]) \sqcup \mathrm{gr}_\bullet(\mathbb{F}_p[\![F]\!]).$$

It is worth observing that by Theorem 5.1 one-relator maximal pro-p Galois groups "behave" like such free pro-p products in terms of $\mathbb{F}_p$-cohomology and restricted Lie algebras. In other words, the algebras $H^\bullet(G)$ and $\mathrm{gr}_\bullet(\mathbb{F}_p[\![G]\!])$ cannot "distinguish" between such free pro-p products and other one-relator maximal pro-p Galois groups.

Example 5.3. In [KZ05], D. Kochloukova and P. Zalesskii produced the following example of free-by-Demushkin pro-p group which is not such a free pro-p product: let G be the pro-p group with minimal presentation

$$G = \langle x_1, x_2, x_3 \mid [x_1, x_2] = x_3^q \rangle ,$$

with $q \geq 0$ a p-power, $q \neq 2$ if $p = 2$. Such pro-p group satisfies all the conditions in Würfel's theorem (cf. [KZ05, Thm. 2]), and also Proposition 4.1. In particular, one has

$$H^\bullet(G) = H^\bullet(\bar{G}) \sqcap H^\bullet(C),$$

with $C = \langle x_3 \rangle$ and $\bar{G} = G/N \simeq \mathbb{Z}_p^2$, with N the normal closure of C, and

$$\mathrm{gr}_\bullet(\mathbb{F}_p[\![G]\!]) \simeq \mathbb{F}_p[X_1, X_2] \sqcup \mathbb{F}_p[X_3].$$

Yet, the group G is not realizable as maximal pro-p Galois group of any field (see also [Qua15, § 4.2.1]).

Acknowledgment. The author wishes to express his gratitude to I. Efrat and J.P. Labute, for their precious comments and suggestions, to J. Mináč and N.D. Tân, for the stimulating time spent working together at the University of Western Ontario, and to F.W. Pasini for the inspiring discussions about quadratic algebras. Also, he owes many thanks to C. Maire, E. Matzri, D. Neftin, I. Snopce and J. Sonn for their interest.

References

[DdSMS99] J. D. Dixon, M. P. F. du Sautoy, A. Mann, and D. Segal, *Analytic pro-p groups*, second ed., Cambridge Studies in Advanced Mathematics, vol. 61, Cambridge University Press, Cambridge, 1999. MR 1720368 (2000m:20039)

[Efr95] Ido Efrat, *Orderings, valuations, and free products of Galois groups*, Sem. Structure Algébriques Ordonnées, Univ. Paris VII (1995).

[Efr98] ———, *Small maximal pro-p Galois groups*, Manuscripta Math. **95** (1998), no. 2, 237–249. MR 1603329 (99e:12005)

[Gär15] Jochen Gärtner, *Higher Massey products in the cohomology of mild pro-p-groups*, J. Algebra **422** (2015), 788–820. MR 3272101

[KZ05] Dessislava H. Kochloukova and Pavel Zalesskii, *Free-by-Demushkin pro-p groups*, Math. Z. **249** (2005), no. 4, 731–739. MR 2126211 (2005j:20030)

[Lab67] John P. Labute, *Classification of Demushkin groups*, Canad. J. Math. **19** (1967), 106–132. MR 0210788 (35 #1674)

[Lab06] ———, *Mild pro-p-groups and Galois groups of p-extensions of* $\mathbb{Q}$, J. Reine Angew. Math. **596** (2006), 155–182. MR 2254811 (2007j:11158)

[LV12] Jean-Louis Loday and Bruno Vallette, *Algebraic operads*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 346, Springer, Heidelberg, 2012. MR 2954392

[MPQT] J. Mináč, F. W. Pasini, C. Quadrelli, and N. D. Tân, *Quadratic duals and koszul algebras in galois cohomology*, preprint.

[MS82] A. S. Merkur'ev and A. A. Suslin, *K-cohomology of Severi-Brauer varieties and the norm residue homomorphism*, Izv. Akad. Nauk SSSR Ser. Mat. **46** (1982), no. 5, 1011–1046, 1135–1136. MR 675529 (84i:12007)

[NSW08] Jürgen Neukirch, Alexander Schmidt, and Kay Wingberg, *Cohomology of number fields*, second ed., Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 323, Springer-Verlag, Berlin, 2008. MR 2392026 (2008m:11223)

[Pos05] Leonid Positselski, *Koszul property and Bogomolov's conjecture*, Int. Math. Res. Not. (2005), no. 31, 1901–1936. MR 2171198 (2006h:19002)

[Pos14] ———, *Galois cohomology of a number field is Koszul*, J. Number Theory **145** (2014), 126–152. MR 3253297

[PP05] Alexander Polishchuk and Leonid Positselski, *Quadratic algebras*, University Lecture Series, vol. 37, American Mathematical Society, Providence, RI, 2005. MR 2177131 (2006f:16043)

[Pri70] Stewart B. Priddy, *Koszul resolutions*, Trans. Amer. Math. Soc. **152** (1970), 39–60. MR 0265437 (42 #346)

[PV95] Leonid Positselski and Alexander Vishik, *Koszul duality and Galois cohomology*, Math. Res. Lett. **2** (1995), no. 6, 771–781. MR 1362968 (97b:12008)

[Qua14] Claudio Quadrelli, *Bloch-Kato pro-p groups and locally powerful groups*, Forum Math. **26** (2014), no. 3, 793–814. MR 3200350

[Qua15] ———, *Cohomology of absolute galois*, Ph.D. thesis, University of Western Ontario, 2015, UWO Electronic Thesis and Dissertation Repository. Paper 2600.

[QW] Claudio Quadrelli and Thomas S. Weigel, *Profinite groups with a cyclotomic orientation*, preprint.

[Wei15a] Thomas Weigel, *Graded Lie algebras of type FP*, Israel J. Math. **205** (2015), no. 1, 185–209. MR 3314587

[Wei15b] ———, *Koszul Lie algebras*, Lie algebras and related topics (M. Avitabile, J. Feldvoss, and Th. Weigel, eds.), Contemporary Mathematics, vol. 652, American Mathematical Society, Providence, RI, 2015, pp. 254–255.

[Wür85] Tilmann Würfel, *A remark on the structure of absolute Galois groups*, Proc. Amer. Math. Soc. **95** (1985), no. 3, 353–356. MR 806069 (87b:12006)

Department of Mathematics, Ben-Gurion University of the Negev, P.O.B. 653, 84105 Be'er-Sheva, Israel

E-mail address: quadrell@math.bgu.ac.il